Amlapitta: A Disease-Focused Narrative Review of Ayurvedic Concepts and Contemporary gastroenterology

Nitin Maroti Ingole[1*], Bhagyesha K[2]

[1]*Dept. of P.G. Studies in Kayachikitsa, K.V.G. Ayurveda Medical College And Hospital, Ambate Adka, Sullia, D. K. District, Karnataka, India-574327.*

[2]*Professor And Hod, Dept. of P.G. Studies In Kayachikitsa, K.V.G. Ayurveda Medical College And Hospital, Ambate Adka, Sullia, D. K. District, Karnataka, India – 574327.*

Corresponding Author:
Nitin Maroti Ingole
E-mail: nitiningole3103@gmail.com

Submission: 29.07.2026 Acceptance: 28.09.2026 Publication: 06.11.2026

https://www.doi.org/10.63778/CIJISM-ARJCPL/2026_11264

Abstract

Amlapitta is described in Ayurveda as a disorder involving Vidagdha Pitta and impaired digestive function (Agnimandya), with manifestations that may include sour eructation, burning discomfort, nausea, reduced appetite, abdominal discomfort and vomiting. Classical descriptions discuss dietary, lifestyle and psychological factors, and classify clinical patterns according to Dosha predominance and direction of manifestation (Urdhwaga and Adhoga). Some symptoms overlap with gastroesophageal reflux disease (GERD), dyspepsia and other acid-related disorders; however, these are not interchangeable diagnoses. GERD concerns troublesome reflux symptoms or complications due to retrograde flow of gastric contents, whereas dyspepsia includes epigastric pain or burning, postprandial fullness and early satiety. This narrative review synthesizes the Amlapitta description in the supplied dissertation with contemporary diagnostic and management principles. It discusses Nidana, Samprapti, clinical features, differential diagnosis, red flags, lifestyle measures and research priorities. Ayurvedic constructs may guide individualized assessment, but their correspondence with biomedical diagnoses requires explicit operational definitions and validation. Persistent or alarm symptoms require appropriate medical evaluation. Better-designed observational studies and controlled trials using validated outcomes are needed to assess Ayurvedic interventions without overstating current evidence.

Keywords: Amlapitta; Ayurveda; Agnimandya; Pitta; gastroesophageal reflux disease; dyspepsia; narrative review

Introduction

Amlapitta is a recognized disease entity in later Ayurvedic literature, particularly in descriptions attributed to Kashyapa, Madhava Nidana and subsequent compendia. The supplied dissertation describes Amlapitta through the concepts of Pitta vitiation, sourness (Amla guna), impaired digestive function and the transformation of ingested food under conditions of Agnimandya. Its symptom profile includes Amlodgara, Hrid-Kantha Daha, Hridshoola, Aruchi, Adhmana, Angasada and, in the study proforma, Shirashoola[1-3]. In contemporary practice, symptoms commonly labelled "hyperacidity" may arise from several conditions, including GERD, functional dyspepsia, peptic ulcer disease, medication-related mucosal injury and other gastrointestinal disorders. GERD is not simply excessive acid production; reflux, esophageal clearance, mucosal integrity and symptom sensitivity also influence presentation[4,5]. A disease-focused review should therefore describe the Ayurvedic framework while distinguishing conceptual overlap from diagnostic equivalence.

Review approach

This narrative review is based primarily on the Amlapitta sections of the supplied dissertation, including its historical, Ayurvedic and modern reviews and its listed classical-text bibliography. Contemporary clinical context was supplemented with established gastroenterology guidance and peer-reviewed literature. This is not a systematic review: no reproducible database search strategy, formal risk-of-bias assessment or quantitative synthesis was undertaken. Ayurvedic disease descriptions are presented as source-derived concepts; biomedical interpretation is distinguished from them.

Ayurvedic disease concept

Definition and textual development

The dissertation summarizes traditional explanations of Amlapitta as a state in which Pitta becomes excessive in its Amla or Vidahi qualities. It reports that Amlapitta is not presented as a distinct disease in Vedic literature and is not consistently treated as a separate entity in the earliest major Samhitas, while more explicit descriptions occur in later texts. Kashyapa Samhita and Madhava Nidana are highlighted for descriptions of clinical features, etiopathogenesis and disease patterns[1–3]. These historical statements should be checked against the specific editions and commentaries before publication, because chapter numbering and interpretations may vary.

Nidana: etiological factors

The source classifies Nidana into dietary (Aharaja), behavioural/lifestyle (Viharaja), psychological (Manasika) and other/exogenous factors. Reported dietary factors include irregular meal timing, eating before the previous meal has been digested, excessive sour, pungent or salty foods, Vidahi foods, incompatible dietary combinations and unsuitable food quantity or preparation. Lifestyle factors include night-waking, excessive exertion, fasting, daytime sleeping and suppression of natural urges. Stress, worry and grief are also discussed. Smoking, alcohol and prolonged NSAID use are mentioned among possible aggravating factors[1].

These factors belong to the Ayurvedic explanatory framework and should not all be presented as individually proven causes. Modern evidence indicates that triggers vary between individuals; obesity, late meals and smoking can aggravate reflux in some patients, while NSAIDs are particularly relevant to mucosal injury and ulcer risk. Clinical history should document actual triggers and medication exposure rather than impose a universal avoidance list[4,5].

Purvarupa and Rupa

The dissertation notes that classical sources do not consistently specify a separate Purvarupa for Amlapitta. Agnimandya and Ajirna are proposed as possible preceding features. Principal symptoms recorded in its clinical assessment are Amlodgara (sour eructation), Hrid-Kantha Daha (burning in the chest/throat region), Hridshoola (pain/discomfort), Aruchi (reduced desire for food), Adhmana (abdominal distension), Angasada (lassitude) and Shirashoola (headache). Other features in the literature review include nausea, vomiting, excessive salivation, bitter or sour taste, heaviness and reduced digestive appetite[1]. These symptoms are clinically nonspecific. Chest pain must not automatically be attributed to Amlapitta or reflux, particularly if severe, exertional or associated with breathlessness, sweating or faintness; urgent assessment may be necessary to exclude cardiac causes.

Classification

The dissertation describes classification by Dosha predominance-Vataja, Pittaja and Kaphaja-and by direction of manifestation as Urdhwaga and Adhoga Amlapitta. Vata-associated features include pain, tremor, dizziness or weakness; Pitta-associated features include burning, sour or bitter eructation, nausea and thirst; Kapha-associated features include heaviness, reduced appetite, salivation and vomiting. Urdhwaga descriptions emphasize upward symptoms such as eructation and vomiting, whereas Adhoga descriptions include lower gastrointestinal or systemic features in the cited classical accounts[1–3].

These are traditional clinical classifications, not validated biomedical subtypes. Research should define the criteria prospectively and report the symptoms used to assign each category.

Samprapti and Samprapti Ghataka

The disease process summarized in the dissertation begins with Nidana that aggravate Doshas, particularly Pitta and Vata, and impair Jatharagni. Food is described as undergoing Vidagdha transformation, with sour/fermented characteristics and persistence of symptoms. Continued Dosha-Dushya interaction is proposed to produce characteristic Amlapitta manifestations, later differentiated into Urdhwaga and Adhoga patterns[1]. This is a traditional pathophysiological model rather than a directly demonstrated biochemical sequence.

For research, Samprapti Ghataka should be reported explicitly rather than inferred after treatment response. Relevant elements may include the proposed Dosha, Dushya, Agni status, Ama concept, Srotas, Udbhava Sthana and Vyakta Sthana, with transparent assessment rules. This would improve reproducibility and comparison across studies.

Relationship with contemporary gastrointestinal disorders

Amlapitta and GERD

GERD is a condition in which reflux of gastric contents causes troublesome symptoms or complications. Typical symptoms include heartburn and regurgitation; extra-esophageal complaints have many alternative causes. Diagnosis may be based on typical symptoms and response to an appropriately administered PPI trial in patients without alarm features, while endoscopy and reflux monitoring are used when indicated[4,5].

Amlodgara and Hrid-Kantha Daha may overlap symptomatically with regurgitation and heartburn, but this does not establish that every person classified as having Amlapitta has GERD. Amlapitta includes a broader Ayurvedic symptom and Dosha framework, whereas GERD has defined biomedical criteria. “Hyperacidity” is imprecise because reflux symptoms can occur without excessive gastric acid secretion.

Amlapitta and dyspepsia

Dyspepsia commonly includes epigastric pain or burning, postprandial fullness and early satiety; nausea, bloating and belching may accompany it. Functional dyspepsia is considered when symptoms are not explained by an identifiable structural disease after appropriate evaluation. GERD and dyspepsia can coexist, and symptom overlap complicates classification[6,7]. Amlapitta symptoms such as Hridshoola, Aruchi and Adhmana may overlap with

dyspepsia, but the mapping is partial and should be handled cautiously.

Differential diagnosis and alarm features

Depending on presentation, differential diagnoses include GERD, functional dyspepsia, gastritis, peptic ulcer disease, medication-associated mucosal injury, biliary disease and non-gastrointestinal causes of chest or upper abdominal pain. Helicobacter pylori testing may be relevant in dyspepsia according to age, local guidance and clinical context. Alarm features-including progressive dysphagia, gastrointestinal bleeding, iron-deficiency anaemia, persistent vomiting, unexplained weight loss or a palpable mass-warrant prompt medical assessment[4,5,8].

Assessment and diagnosis

An Ayurvedic assessment may document Nidana, Dosha predominance, Agni, bowel habits, appetite, symptom direction and the classical Lakshana profile. Research should standardize symptom definitions, use validated scales where available, and assess outcomes at prespecified time points. Baseline data should include symptom duration and frequency, nocturnal symptoms, medication use, diet, comorbidities and prior investigations.

Biomedical assessment should be guided by the suspected disorder and risk profile. Typical reflux symptoms without alarm features may be managed initially according to guidelines; endoscopy is indicated in selected patients, including those with alarm symptoms or persistent symptoms despite appropriate management. Reflux monitoring and manometry are reserved for specific diagnostic questions [4,5]. Ayurvedic classification should complement-not delay-appropriate investigation or referral.

Management principles

Ayurvedic approach

The source describes three broad principles: Nidana Parivarjana (avoidance or modification of causative factors), Prakritivighata/Shamana (management directed at the proposed Dosha disturbance) and Apakarshana (elimination procedures selected according to clinical context). It discusses dietary and lifestyle regulation and Shamana measures with properties traditionally considered Pitta-pacifying. Vamana, Virechana or Basti are mentioned in classical discussions according to the type and site of Dosha involvement[1–3].

These interventions should not be presented as universally indicated or as established equivalents to GERD treatment. Shodhana procedures require qualified assessment, an appropriate setting, informed consent and consideration of contraindications. The evidence and safety profile of each intervention must be assessed separately.

Diet and lifestyle

The dissertation recommends regular meals, lighter food and avoidance of foods perceived to aggravate symptoms, including excessively spicy, sour or salty items. Contemporary GERD guidance supports individualized trigger-food avoidance, weight reduction for people with overweight or obesity, avoiding meals close to bedtime, head-of-bed elevation for nocturnal symptoms and smoking cessation where relevant. Evidence does not support imposing every dietary restriction on every patient; advice should be individualized to observed triggers and nutritional needs[4,5].

Contemporary medical care and integrative practice

Antacids or alginate-based products may provide short-term relief in selected patients, while PPIs are principal pharmacological treatments for many patients with GERD, particularly erosive disease. Indication, dose and duration should be clinically appropriate, with reassessment of response. Persistent symptoms should prompt review of adherence, medication timing, diagnosis and alternative causes rather than indefinite escalation without reassessment [4,5].

Where Ayurvedic care is considered, it should be integrated with appropriate diagnostic assessment and medication review. Product identity, botanical authentication, preparation standards, dose, duration, adverse events and concomitant medicines should be documented. Patients should not discontinue prescribed medicines or delay evaluation of alarm symptoms because of an Ayurvedic disease label.

Evidence gaps and research priorities

The supplied dissertation reports an open, single-group clinical study of 30 adults treated with Sa Chana Saktu Panchanimbadi Churna for 14 days, with a further seven-day follow-up. It reports symptom-score improvement and no observed adverse effects during the short observation period. Because there was no control group, blinding or longer follow-up, these findings are preliminary and cannot establish comparative efficacy, causal attribution or long-term safety. They concern one formulation and should not be interpreted as evidence for all Amlapitta treatments[1].

Future research should define Amlapitta inclusion criteria and Urdhwaga/Adhoga classifications prospectively; distinguish probable GERD, dyspepsia and other diagnoses; use validated symptom and quality-of-life instruments; prespecify outcomes and sample-size calculations; use suitable control groups and blinded outcome assessment where feasible; standardize interventions and co-interventions; systematically record adverse events; and

include adequate follow-up. Pragmatic trials could assess whether individualized Ayurvedic assessment adds measurable value to guideline-based care.

Limitations of this review

This is a narrative review, not a systematic review. The Ayurvedic disease description is derived chiefly from the supplied dissertation and its listed classical references; exact Sanskrit passages and chapter/verse details were not independently collated for this manuscript. The biomedical discussion uses selected guidelines and reviews rather than an exhaustive evidence search. This article therefore provides a conceptual synthesis and research framework, not a definitive statement of treatment efficacy.

Conclusion

Amlapitta is described in Ayurveda through the linked concepts of Pitta vitiation, Agnimandya, dietary and behavioural Nidana, and symptom patterns classified by Dosha predominance and direction of manifestation. Several symptoms overlap with GERD and dyspepsia, but the disorders should not be treated as synonymous. A clinically responsible approach combines Ayurvedic assessment with biomedical differential diagnosis, recognition of alarm features and individualized lifestyle advice. Evidence for specific Ayurvedic interventions remains preliminary when studies are small and uncontrolled. Standardized disease definitions and well-designed comparative research are essential before strong claims can be made about efficacy or safety.

Declarations

Ethics approval: Not applicable

Data availability: Not applicable.

Source of Support: Nil

Conflict of Interest: Nil

Copyright © 2026 CSMSS International Journal of Indian System of Medicine. This is an open access article, it is free for all to read, download, copy, distribute, adapt and permitted to reuse under Creative Commons Attribution Non Commercial-ShareAlike: CC BY-NC-SA BY 4.0 license.

References

1. Author and bibliographic details to be confirmed. A Comprehensive Clinical Study of Sa Chana Saktu Panchanimbadi Churna in Amlapitta [dissertation]. Sullia: KVG Ayurveda Medical College; year to be confirmed. (Source dissertation supplied for this review.)
2. Madhavakara. Madhava Nidana with Madhukosha Commentary. Upadhyaya Y, editor. Varanasi: Chaukhambha Prakashan; 2019.
3. Vriddha Jivaka. Kashyapa Samhita. Sharma H, editor. Varanasi: Chaukhambha Sanskrit Sansthan; 2018.
4. Katz PO, Dunbar KB, Schnoll-Sussman FH, Greer KB, Yadlapati R, Spechler SJ. ACG Clinical Guideline for the Diagnosis and Management of Gastroesophageal Reflux Disease. Am J Gastroenterol. 2022;117(1):27-56. doi:10.14309/ajg.0000000000001538.
5. Yadlapati R, Gyawali CP, Pandolfino JE, et al. AGA Clinical Practice Update on the Personalized Approach to the Evaluation and Management of GERD: Expert Review. Clin Gastroenterol Hepatol. 2022;20(5):984-994.e1. doi:10.1016/j.cgh.2022.01.025.
6. Drossman DA. Functional Gastrointestinal Disorders: History, Pathophysiology, Clinical Features and Rome IV. Gastroenterology. 2016;150(6):1262-1279. doi:10.1053/j.gastro.2016.02.032.
7. Harer KN, Hasler WL. Functional Dyspepsia: A Review of the Symptoms, Evaluation, and Treatment Options. Gastroenterol Hepatol (N Y). 2021;17(2):84-91. Verify final bibliographic details before submission.
8. World Gastroenterology Organisation. Gastroesophageal Reflux Disease (GERD): Global Guideline [Internet]. Available from: https://www.worldgastroenterology.org/guidelines / gastroesophageal-reflux-disease. Accessed 18 October 2026.
9. World Gastroenterology Organisation. Common GI Symptoms: Practice Guideline [Internet]. Available from: https://www.worldgastroenterology.org/guidelines/common-gi-symptoms. Accessed 18 October 2026.
10. Agnivesha. Charaka Samhita. Revised by Charaka and Dridhabala. Shastri K, Chaturvedi G, editors. Varanasi: Chaukhambha Bharati Academy; 2020.
11. Sushruta. Sushruta Samhita. Shastri AD, editor. Varanasi: Chaukhambha Sanskrit Sansthan; 2021.
12. Govinda Das Sen. Bhaishajya Ratnavali. Shastri AD, editor. Varanasi: Chaukhambha Prakashan; 2020.